

A new two-dimensional lattice model that is “consistent around a cube”

Jarmo Hietarinta

Department of Physics, University of Turku

FIN-20014 Turku, FINLAND

e-mail: Jarmo.Hietarinta@utu.fi

February 8, 2008

Abstract

For two-dimensional lattice equations one definition of integrability is that the model can be naturally and consistently extended to three dimensions, i.e., that it is “consistent around a cube” (CAC). As a consequence of CAC one can construct a Lax pair for the model. Recently Adler, Bobenko and Suris conducted a search based on this principle and certain additional assumptions. One of those assumptions was the “tetrahedron property”, which is satisfied by most known equations. We present here one lattice equation that satisfies the consistency condition but does not have the tetrahedron property. Its Lax pair is also presented and some basic properties discussed.

1 Introduction

Within the field of integrable dynamics there have recently been interesting developments in the study of integrable *difference* equations (an overview of the topic can be obtained from the proceedings mentioned in [1]). These include both the discussion on what is a proper definition of integrability, or whether the different suggested definitions actually agree or not. Numerous integrable (in some sense) difference equations have been proposed by discretizing known ODE’s and PDE’s in a way that retains some good properties. Once a definition of integrability has been proposed one can also try to search for all equations having the chosen property.

In this paper we consider integrable difference equations defined on a 2-dimensional lattice. We assume that the lattice is rectangular and infinite, and concentrate on some square in it (see Figure 1) (We normally use subscripts in square brackets to indicate shifts in the indices, but for composite expression we sometimes also use $\sim$ for shift in the 1-direction and $\hat{\cdot}$ for a shift in the 2-direction.) In general the map is given in terms of a multi-linear equation relating the four corner values in Figure 1:

$$\begin{aligned} & K xx_{[1]x_{[2]}x_{[12]}} + L_1 xx_{[1]x_{[2]}} + L_2 xx_{[1]x_{[12]}} + L_3 xx_{[2]x_{[12]}} + L_4 x_{[1]x_{[2]}x_{[12]}} \\ & + P_1 xx_{[1]} + P_2 x_{[1]x_{[2]}} + P_3 x_{[2]x_{[12]}} + P_4 x_{[12]x} + P_5 xx_{[2]} + P_6 x_{[1]x_{[12]}} \\ & + Q_1 x + Q_2 x_{[1]} + Q_3 x_{[2]} + Q_4 x_{[12]} + U \equiv Q_{12}(x, x_{[1]}, x_{[2]}, x_{[12]}; p_1, p_2) = 0. \end{aligned} \quad (1)$$




Figure 1: The lattice map is defined on a elementary square of the lattice

Here the coefficients $K, L_\nu, P_\nu, Q_\nu, U$ may depend on the two spectral parameters p_1, p_2 . If any 3 of the corner values are given then the fourth one can be obtained as a rational expression of the other three. One can therefore propagate any staircase-like initial value line to cover the whole plane[2].

How should integrability be defined for such maps? In [3] the following definition of integrability (“Consistency Around the Cube”, CAC) was proposed (see Figure 2): Adjoin a third direction (therefore assuming $x = x_{n,m,k}$) and use the same map (but with different spectral parameters) also in planes corresponding to indices 1,3 and 2,3. That is, the map given in (1) contains shifts and parameters associated with directions 1,2 now the same should be done with directions 3,1 and 2,3, furthermore, on the parallel shifted planes we use identical maps. We assume that the values $x, x_{[1]}, x_{[2]}, x_{[3]}$ at black circles in Figure 2 are given, then the values at open circles are uniquely determined using the relevant map, but the value at $x_{[123]}$ can be computed in 3 different ways, and they must give the same result. In other words:

$$\begin{aligned} \text{solve } x_{[12]} \text{ from } & Q_{12}(x, x_{[1]}, x_{[2]}, x_{[12]}; p_1, p_2) = 0, \\ \text{solve } x_{[23]} \text{ from } & Q_{23}(x, x_{[2]}, x_{[3]}, x_{[23]}; p_2, p_3) = 0, \\ \text{solve } x_{[31]} \text{ from } & Q_{31}(x, x_{[3]}, x_{[1]}, x_{[31]}; p_3, p_1) = 0, \end{aligned}$$




Figure 2: Given the values at the black circles, one should get a unique value for $x_{[123]}$, even though there are three possible way to compute it.

then $x_{[123]}$ computed from

$$\begin{aligned} Q_{12}(x_{[3]}, x_{[31]}, x_{[23]}, x_{[123]}; p_1, p_2) &= 0, & \text{or} \\ Q_{23}(x_{[1]}, x_{[12]}, x_{[31]}, x_{[123]}; p_2, p_3) &= 0, & \text{or} \\ Q_{31}(x_{[2]}, x_{[23]}, x_{[12]}, x_{[123]}; p_3, p_1) &= 0, \end{aligned}$$

should be the same. The functions Q_{ij} could be in principle be different, but are usually assumed to be identical.

The above idea (and diagram) resembles, e.g., the 3D Bianchi diagram that is obtained from consistency of Moutard transformations (see [4]), but it is here used in a different context: the diagram introduces a condition for lattice maps defined on *squares* rather than for Moutard transformations defined on *edges*. Furthermore, this CAC principle is a *constructive* definition of integrability in the sense that it leads algorithmically to a Lax pair.

The following maps are well known examples that have CAC property[1, 2]:

1. Lattice KdV: $(p_1 - p_2 + x_{[2]} - x_{[1]})(p_1 + p_2 + x - x_{[12]}) = p_1^2 - p_2^2$,
2. Lattice MKdV: $p_1(xx_{[2]} - x_{[1]}x_{[12]}) = p_2(xx_{[1]} - x_{[2]}x_{[12]})$,
3. Lattice SKdV: $(x - x_{[2]})(x_{[1]} - x_{[12]})p_2^2 = (x - x_{[1]})(x_{[2]} - x_{[12]})p_1^2$.

2 Searching for integrable lattices

Now that the definition of integrability has been given one may ask for a listing of all integrable models. The complete classification of CAC maps is in fact a formidable open problem. The set of equations that needs to be solved can be derived from above: comparing the different forms for $x_{[123]}$ and collecting the various coefficients of x and its shifts, we get 2×375 functional equations for the coefficient functions (in practice we have often used 3×375 equations for a symmetric approach). The equations are polynomial in the coefficient functions appearing in (1), but the functions depend on different pairs of the three spectral parameters λ_i , $i = 1, 2, 3$ and all three λ_i 's appear in each equation. The equation list itself takes 35MB to store.

In [5] the consistency equations were solved in the case where the map Q is the same on all planes, and under two additional assumptions:

- symmetry:

$$\begin{aligned} Q(x, x_{[1]}, x_{[2]}, x_{[12]}; p_1, p_2) &= \varepsilon Q(x, x_{[2]}, x_{[1]}, x_{[12]}; p_2, p_1) \\ &= \sigma Q(x_{[1]}, x, x_{[12]}, x_{[2]}; p_1, p_2), \quad \varepsilon, \sigma = \pm 1, \end{aligned} \tag{2}$$

- “tetrahedron property”: $x_{[123]}$ does not depend on x .

With these assumptions the authors were able to get a full classification resulting with 9 models.

The tetrahedron assumption is indeed satisfied by most of the well known models, but one can nevertheless ask whether it is a fundamental or essential property and whether

there are any nontrivial models that do not satisfy it. It should be immediately observed that the more or less trivial models

$$x_{[1]}x_{[2]} - xx_{[12]} = 0, \text{ and } y - y_{[1]} - y_{[2]} + y_{[12]} = 0, \quad (3)$$

($x = e^y$) do have CAC property but *not* the tetrahedron property, in fact one quickly finds that

$$x_{[123]} = x_{[1]}x_{[2]}x_{[3]}x^{-2}, \text{ and } y_{[123]} = y_{[1]} + y_{[2]} + y_{[3]} - 2y, \quad (4)$$

respectively. One may now ask whether these models have nontrivial extensions.

The symmetry assumptions (2) lead to 2 possibilities w.r.t. σ , the one with $\sigma = -1$ is:

$$\begin{aligned} Q = & a_1(p_1, p_2)(xx_{[1]}x_{[2]} - xx_{[2]}x_{[12]} - xx_{[1]}x_{[12]} + x_{[1]}x_{[2]}x_{[12]}) \\ & + a_2(p_1, p_2)(xx_{[12]} - x_{[1]}x_{[2]}) + a_3(p_1, p_2)(x - x_{[1]} - x_{[2]} + x_{[12]}). \end{aligned} \quad (5)$$

In [5] it was observed that in this symmetry class there are no integrable cases with tetrahedron property, which is true. The ansatz (5) forms, fortunately, a rather simple class and a direct computation shows that it contains one new integrable map

$$\begin{aligned} & xx_{[1]}x_{[2]} - xx_{[1]}x_{[12]} - xx_{[12]}x_{[2]} + x_{[1]}x_{[2]}x_{[12]} \\ & + (x_{[1]}x_{[2]} - xx_{[12]})(p_1 + p_2) - p_1p_2(x - x_{[1]} - x_{[2]} + x_{[12]}) = 0, \end{aligned} \quad (6)$$

with two parameters. Since it can also be written as

$$\begin{array}{ccccccc} x & x_{[1]} & x_{[2]} & & -pq & x_{[12]} & \\ -x & x_{[1]} & x_{[12]} & + (p+q) & x_{[1]} & x_{[2]} & +pq & x_{[2]} \\ -x & x_{[2]} & x_{[12]} & - (p+q) & x & x_{[12]} & +pq & x_{[1]} \\ +x_{[1]} & x_{[2]} & x_{[12]} & & -pq & x & & \end{array} = 0,$$

we call it “the bow-tie model”. Model (6) is, however, still Möbius-equivalent to the first model in (3) by $x \rightarrow -(p_1x + p_2)/(x + 1)$ [although this transformation does not trivialize the maps on the other planes, since it depends explicitly on p_1, p_2]. Therefore we decided to search for possible extensions. This was done perturbatively, with completely general 1st and 2nd order extensions to the bow-tie map (this work was done with REDUCE 3.7[6], and 1GB memory). In this approach it is not necessary to specify the details of the parameter dependence – indeed the result was surprising in this respect. From the outcome of this exercise we noticed certain weaker symmetry properties among the coefficient functions and when the (nonperturbative) computations were done with these properties (and translational invariance) the following solution was found:

$$\begin{aligned} Q(x, x_{[1]}, x_{[2]}, x_{[12]}; e_1, o_1; e_2, o_2) \equiv & \\ & xx_{[1]}x_{[2]}(o_1 - o_2) - xx_{[1]}x_{[12]}(e_1 - o_2) - xx_{[2]}x_{[12]}(o_1 - e_2) + x_{[1]}x_{[2]}x_{[12]}(e_1 - e_2) \\ & + (x_{[1]}x_{[2]} - xx_{[12]})(e_1o_1 - e_2o_2) \\ & + (xx_{[2]}o_1 - x_{[1]}x_{[12]}e_1)(e_2 - o_2) + (x_{[2]}x_{[12]}e_2 - xx_{[1]}o_2)(e_1 - o_1) \\ & - x_{[12]}e_1e_2(o_1 - o_2) + x_{[2]}e_2o_1(e_1 - o_2) + x_{[1]}e_1o_2(o_1 - e_2) - xo_1o_2(e_1 - e_2) \\ & = 0. \end{aligned} \quad (7)$$

This new model is the main result of this paper. It is interesting to note that in (7) there are *two* parameters o_i, e_i in each direction, and that if $e_i = o_i$ then $o_1 - o_2$ factors out

leaving (6). The parameters with different indices must all be different, otherwise the map factorizes. This model obeys the CAC property, in fact one finds

$$\begin{aligned}
\text{num}(x_{[123]}) &= -x(x_{[1]} + o_1)(x_{[2]} + o_2)(x_{[3]} + o_3)(o_1 - o_2)(o_2 - o_3)(o_3 - o_1) \\
&\quad + (x_{[1]} + o_1)(x_{[2]} + o_2)(x_{[3]} + o_3) \\
&\quad \times [(e_1 e_2 + e_3 o_3) o_3 (o_1 - o_2) + (e_2 e_3 + e_1 o_1) o_1 (o_2 - o_3) + (e_3 e_1 + e_2 o_2) o_2 (o_3 - o_1)] \\
&\quad + (x + e_3)(x_{[1]} + o_1)(x_{[2]} + o_2) o_3 (o_1 - o_2)(e_2 - o_3)(o_3 - e_1) \\
&\quad + (x + e_1)(x_{[2]} + o_2)(x_{[3]} + o_3) o_1 (o_2 - o_3)(e_3 - o_1)(o_1 - e_2) \\
&\quad + (x + e_2)(x_{[3]} + o_3)(x_{[1]} + o_1) o_2 (o_3 - o_1)(e_1 - o_2)(o_2 - e_3), \\
\text{den}(x_{[123]}) &= (x_{[1]} + o_1)(x_{[2]} + o_2)(x_{[3]} + o_3) \\
&\quad \times [(e_1 e_2 + e_3 o_3)(o_2 - o_1) + (e_2 e_3 + e_1 o_1)(o_3 - o_2) + (e_3 e_1 + e_2 o_2)(o_1 - o_3)] \\
&\quad + (x + e_3)(x_{[1]} + o_1)(x_{[2]} + o_2)(o_1 - o_2)(e_1 - o_3)(e_2 - o_3) \\
&\quad + (x + e_1)(x_{[2]} + o_2)(x_{[3]} + o_3)(o_2 - o_3)(e_2 - o_1)(e_3 - o_1) \\
&\quad + (x + e_2)(x_{[3]} + o_3)(x_{[1]} + o_1)(o_3 - o_1)(e_3 - o_2)(e_1 - o_2),
\end{aligned}$$

which are symmetric under permutations of 1, 2, 3, and the explicit x -dependence demonstrates violation of the tetrahedron property of [5].

3 Symmetries

The model (7) is invariant under the Möbius transformation

$$\begin{aligned}
X &\rightarrow \frac{\alpha X + \beta}{\gamma X + \delta}, & X = x, x_i, x_{ij}, x_{ijk} \\
P &\rightarrow -\frac{\alpha P - \beta}{\gamma P - \delta}, & P = e_i \text{ or } o_i,
\end{aligned}$$

It has the symmetries

$$Q(x, x_{[1]}, x_{[2]}, x_{[12]}; e_1, o_1, e_2, o_2) = -Q(x, x_{[2]}, x_{[1]}, x_{[12]}; e_2, o_2, e_1, o_1), \quad (8)$$

$$Q(x, x_{[1]}, x_{[2]}, x_{[12]}; e_1, o_1, e_2, o_2) = -Q(x_{[1]}, x, x_{[12]}, x_{[2]}; o_1, e_1, e_2, o_2). \quad (9)$$

These are similar to (2) with $\varepsilon = \sigma = -1$, but note that in (9) there is an exchange between the two parameters related to direction 1. Indeed, it seems that the extension of the parameter space from 1 to 2 dimensional (in each direction of the lattice) allows the introduction of a nontrivial exchange symmetry, resulting with the nontrivial model. (Recall that if $e_i = o_i$ the model simplifies to (6).)

The map (7) can also be written in the following rational form

$$\frac{x + e_2}{x + e_1} \frac{x_{[12]} + o_2}{x_{[12]} + o_1} = \frac{x_{[1]} + e_2}{x_{[1]} + o_1} \frac{x_{[2]} + o_2}{x_{[2]} + e_1}. \quad (10)$$

(The Bäcklund equation presented in Eq. (5.11) of [7] is similar, but the distribution of the terms is in fact different.) A form equivalent to (10) is

$$\frac{o_2 + x_{[12]}}{o_2 + x_{[2]}} \frac{e_2 + x}{e_2 + x_{[1]}} = \frac{o_1 + x_{[12]}}{o_1 + x_{[1]}} \frac{e_1 + x}{e_1 + x_{[2]}},$$

and comparing these one observes duality under

$$o_2 \leftrightarrow x, e_2 \leftrightarrow x_{[12]}, o_1 \leftrightarrow x_{[1]}, e_1 \leftrightarrow x_{[2]}. \quad (11)$$

We also note that (7) has the simple solution

$$x_{n,m,p} = c + (e_1 - o_1)n + (e_2 - o_2)m + (e_3 - o_3)p.$$

4 Lax pair

An important property of the CAC definition of integrability is that it provides a method of constructing a Lax pair. The recipe was provided by Nijhoff in [8]: We consider the three maps on planes 12, 23, 31. The third direction is taken as auxiliary (spectral) and the system is linearized in the corresponding variable $x_{[3]}$ and its shifts. We also use notation $o_3 = \mu$, $e_3 = \lambda$.

Solving for $x_{[31]}$ from $Q(x, x_{[3]}, x_{[1]}, x_{[31]}; \lambda, \mu, e_1, o_1) = 0$ yields

$$x_{[31]} = \frac{x_{[3]}[xx_{[1]}(o_1 - \mu) + xo_1(\lambda - \mu) + x_{[1]}(e_1 o_1 - \lambda \mu) + \lambda o_1(e_1 - \mu)] + xx_{[1]}\mu(-e_1 + o_1) + xo_1\mu(-e_1 + \lambda) + x_{[1]}e_1\mu(-\lambda + o_1)}{x_{[3]}[x_{[1]}(-e_1 + \lambda) + x(-\lambda + o_1) + \lambda(-e_1 + o_1)] + xx_{[1]}(e_1 - \mu) + x(e_1 o_1 - \lambda \mu) + x_{[1]}e_1(\lambda - \mu) + e_1\lambda(o_1 - \mu)}$$

and this is then linearized by introducing f, g by

$$x_{[3]} = \frac{f}{g}, \quad x_{[23]} = \frac{f_{[2]}}{g_{[2]}}, \quad x_{[31]} = \frac{f_{[1]}}{g_{[1]}}, \quad (12)$$

resulting in

$$\begin{aligned} f_{[1]} &= \kappa_1 f [xx_{[1]}(\mu - o_1) + xo_1(\mu - \lambda) + x_{[1]}(\lambda\mu - e_1 o_1) + \lambda o_1(\mu - e_1)] \\ &\quad + \kappa_1 g [xx_{[1]}\mu(e_1 - o_1) + xo_1\mu(e_1 - \lambda) + x_{[1]}e_1\mu(\lambda - o_1)], \\ g_{[1]} &= \kappa_1 f [x_{[1]}(e_1 - \lambda) + x(\lambda - o_1) + \lambda(e_1 - o_1)] \\ &\quad + \kappa_1 g [xx_{[1]}(\mu - e_1) + x(\lambda\mu - e_1 o_1) + x_{[1]}e_1(\mu - \lambda) + e_1\lambda(\mu - o_1)]. \end{aligned}$$

Here the overall separation factor κ_i may contain $x, x_{[i]}$. We write this, and the corresponding equation obtained from $Q(x, x_{[2]}, x_{[3]}, x_{[23]}; e_2, o_2, \lambda, \mu) = 0$ in matrix form,

$$\phi = \begin{pmatrix} f \\ g \end{pmatrix}, \quad \phi_{[i]} = \begin{pmatrix} f_{[i]} \\ g_{[i]} \end{pmatrix}, \quad \phi_{[i]} = L_i \phi, \quad (13)$$

with

$$L_i = \kappa_i(x_{[i]} + o_i)(x + \lambda) \begin{pmatrix} \mu & \mu e_i \\ -1 & -e_i \end{pmatrix} + \kappa_i(x_{[i]} + \lambda)(x + e_i) \begin{pmatrix} -o_i & -\mu o_i \\ 1 & \mu \end{pmatrix}. \quad (14)$$

Now from the consistency condition $(\phi_{[1]})_{[2]} = (\phi_{[2]})_{[1]}$ we get the matrix equation

$$\hat{L}_1 L_2 = \tilde{L}_2 L_1, \quad (15)$$

which is satisfied modulo the map (7), provided that the separation constants κ_i satisfy

$$\frac{\kappa_1 \tilde{\kappa}_2}{\widehat{\kappa}_1 \kappa_2} = \frac{(x_{[2]} + e_1)(x + e_2)(x_{[2]} + \lambda)}{(x + e_1)(x_{[1]} + e_2)(x_{[1]} + \lambda)}. \quad (16)$$

A simple solution to this is

$$\kappa_i = \frac{1}{(x + e_i)(x + \lambda)}, \quad (17)$$

leading to

$$L_i = \frac{x_{[i]} + o_i}{x + e_i} \begin{pmatrix} \mu & \mu e_i \\ -1 & -e_i \end{pmatrix} + \frac{x_{[i]} + \lambda}{x + \lambda} \begin{pmatrix} -o_i & -\mu o_i \\ 1 & \mu \end{pmatrix} \quad (18)$$

A similarity transformation with $S = \begin{pmatrix} 1 & 0 \\ \mu^{-1} & 1 \end{pmatrix}$ simplifies this further to

$$L'_i = \begin{pmatrix} (\mu - e_i) \frac{x_{[i]} + o_i}{x + e_i} & \mu e_i \frac{x_{[i]} + o_i}{x + e_i} - \mu o_i \frac{x_{[i]} + \lambda}{x + \lambda} \\ 0 & (\mu - o_i) \frac{x_{[i]} + \lambda}{x + \lambda} \end{pmatrix}. \quad (19)$$

5 Summary

The “consistency around the cube” definition of integrability is very transparent since it directly leads to the Lax pair. It also defines a clear search problem whose complete solution is, unfortunately, still beyond our computational abilities. The integrable model (7) or (10) presented in this paper is the first result in our search project. This model has several interesting features:

- It does *not* satisfy the tetrahedron property.
- It has *two* parameters in each direction.
- It is dual under interchange of variables and parameters.

Further properties of this model will be discussed elsewhere.

In this general class of lattice models (multi-linear, consistent around a cube) there are probably still many other models to be found.

Acknowledgements

The author would like to thank Frank Nijhoff for numerous discussions.

References

- [1] “Symmetries and Integrability of Difference Equations”, eds: D Levi, L Vinet and P Winternitz, CRM Proceedings & Lecture Notes, vol. 9, (American Mathematical Society, 1996); “Symmetries and Integrability of Difference Equations”, eds: PA Clarkson, and FW Nijhoff, LMS Lecture Note Series, vol. 255, (Cambridge U P, 1999); “SIDE III–Symmetries and Integrability of Difference Equations”, eds: D Levi, and

- O Ragnisco, CRM Proceedings & Lecture Notes, vol. 25, (American Mathematical Society, 2000); “Symmetries and Integrability of Difference Equations: Special Issue Dedicated to the Subject of the SIDE IV Meeting, 27 November–1 December 2000, Tokyo, Japan”, eds: J Hietarinta, F W Nijhoff and J Satsuma, *J. Phys. A.: Math. Gen.* **34** (2001) Number 48.
- [2] FW Nijhoff and HW Capel, *Acta Appl. Math.* **39** (1995) 133.
 - [3] FW Nijhoff and AJ Walker, *Glasgow Math. J* **43A** (2001) 109.
 - [4] JJC Nimmo and WK Schief, *Proc. R. Soc. Lond. A* **453** (1997) 255.
 - [5] V Adler, A Bobenko and Yu Suris, *Commun. Math. Phys.* **233** (2003) 513.
 - [6] AC Hearn: “REDUCE User’s and Contributed Packages Manual”, Version 3.7 (Konrad-Zuse-Zentrum Berlin, Germany, 1999).
 - [7] JJC Nimmo and WK Schief, *Studies in Appl. Math.* **100** (1998) 295.
 - [8] FW Nijhoff, *Phys. Lett. A* **297** (2002) 49.